

Expectation Maximization Mixture Models HMMs

Class 9. 21 Sep 2010


Administrivia

- Homework due Tuesday.
 - Each day of delay thereafter will lose you points
- HW2 will be put up on Tuesday.

A Strange Observation

■ A trend

The pitch of female Indian playback singers is on an ever-increasing trajectory



- Mean pitch values: 278Hz, 410Hz, 580Hz


I'm not the only one to find the high-pitched stuff annoying

- Sarah McDonald (Holy Cow): “.. shrieking...”
- Khazana.com: “.. female Indian movie playback singers who can produce ultra high frequencies which only dogs can hear clearly..”
- www.roadjunky.com: “.. High pitched female singers doing their best to sound like they were seven years old ..”

A Disturbing Observation

■ A trend

The pitch of female Indian playback singers is on an ever-increasing trajectory




- Mean pitch values: 278Hz, 410Hz, 580Hz

A Disturbing Observation

■ A trend

The pitch of female Indian playback singers is on an ever-increasing trajectory



- Mean pitch values: 278Hz, 410Hz, 580Hz

Lets Fix the Song

- The pitch is unpleasant
- The melody isn't bad
- Modify the pitch, but retain melody
- Problem:
 - Cannot just shift the pitch: will destroy the music
 - The music is fine, leave it alone
 - Modify the singing pitch without affecting the music

22 Sep 2011

11755/18797

7

“Personalizing” the Song


- Separate the vocals from the background music
 - Modify the separated vocals, keep music unchanged
- Separation need not be perfect
 - Must only be sufficient to enable pitch modification of vocals
 - Pitch modification is tolerant of low-level artifacts
 - For octave level pitch modification artifacts can be undetectable.

22 Sep 2011

11755/18797

8

Separation example




22 Sep 2011

11755/18797

9

Some examples




- Example 1: Vocals shifted down by 4 semitones

22 Sep 2011

11755/18797

10

Some examples



- Example 1: Vocals shifted down by 4 semitones
- Example 2: Gender of singer partially modified

22 Sep 2011

11755/18797

11

Techniques Employed

- Signal separation
 - Employed a simple latent-variable based separation method
- Voice modification
 - Equally simple techniques
- Will consider the underlying methods over next few lectures
- Separation: Extensive use of Expectation Maximization

22 Sep 2011

11755/18797

12

Learning Distributions for Data

- Problem: Given a collection of examples from some data, estimate its distribution
 - Basic ideas of Maximum Likelihood and MAP estimation can be found in Aarti/Paris' slides
 - Pointed to in a previous class
- Solution: Assign a model to the distribution
 - Learn parameters of model from data
- Models can be arbitrarily complex
 - Mixture densities, Hierarchical models.
- Learning must be done using Expectation Maximization
- Following slides: An intuitive explanation using a simple example of multinomials

22 Sep 2011

11755/18797

13

A Thought Experiment



6 3 1 5 4 1 2 4 ...

- A person shoots a loaded dice repeatedly
- You observe the series of outcomes
- **You can form a good idea of how the dice is loaded**
 - Figure out what the probabilities of the various numbers are for dice
- $P(\text{number}) = \text{count}(\text{number})/\text{sum}(\text{rolls})$
- This is a *maximum likelihood estimate*
 - Estimate that makes the observed sequence of numbers most probable

22 Sep 2011

11755/18797

14

The Multinomial Distribution

- A probability distribution over a discrete collection of items is a *Multinomial*

$$P(X : X \text{ belongs to a discrete set}) = P(X)$$


- E.g. the roll of dice
 - $X : X \text{ in } (1,2,3,4,5,6)$
- Or the toss of a coin
 - $X : X \text{ in } (\text{head}, \text{tails})$

22 Sep 2011

11755/18797

15

Maximum Likelihood Estimation



- Basic principle: Assign a form to the distribution
 - E.g. a multinomial
 - Or a Gaussian
- Find the *distribution* that best fits the histogram of the data

22 Sep 2011

11755/18797

16

Defining "Best Fit"

- The data are generated by draws from the distribution
 - I.e. the generating process draws from the distribution
- Assumption: The distribution has a high probability of generating the observed data
 - Not necessarily true
- Select the distribution that has the *highest* probability of generating the data
 - Should assign lower probability to less frequent observations and vice versa

22 Sep 2011

11755/18797

17

Maximum Likelihood Estimation: Multinomial

- Probability of generating $(n_1, n_2, n_3, n_4, n_5, n_6)$

$$P(n_1, n_2, n_3, n_4, n_5, n_6) = \text{Const} \prod_i p_i^{n_i}$$

- Find $p_1, p_2, p_3, p_4, p_5, p_6$ so that the above is maximized
- Alternately maximize

$$\log(P(n_1, n_2, n_3, n_4, n_5, n_6)) = \log(\text{Const}) + \sum_i n_i \log(p_i)$$

- $\log()$ is a monotonic function
 - $\text{argmax}_x f(x) = \text{argmax}_x \log(f(x))$

- Solving for the probabilities gives us
 - Requires constrained optimization to ensure probabilities sum to 1

$$p_i = \frac{n_i}{\sum_j n_j}$$


**EVENTUALLY
IT'S JUST
COUNTING!**

22 Sep 2011

11755/18797

18

Segue: Gaussians



$$P(X) = N(X; \mu, \Theta) = \frac{1}{\sqrt{(2\pi)^d |\Theta|}} \exp(-0.5(X - \mu)^T \Theta^{-1} (X - \mu))$$

- Parameters of a Gaussian:
 - Mean μ , Covariance Θ

22 Sep 2011

11755/18797

19

Maximum Likelihood: Gaussian

- Given a collection of observations (X_1, X_2, \dots) , estimate mean μ and covariance Θ

$$P(X_1, X_2, \dots) = \prod_i \frac{1}{\sqrt{(2\pi)^d |\Theta|}} \exp(-0.5(X_i - \mu)^T \Theta^{-1} (X_i - \mu))$$

$$\log(P(X_1, X_2, \dots)) = C - 0.5 \sum_i (\log(|\Theta|) + (X_i - \mu)^T \Theta^{-1} (X_i - \mu))$$

- Maximizing w.r.t μ and Θ gives us

$$\mu = \frac{1}{N} \sum_i X_i \quad \Theta = \frac{1}{N} \sum_i (X_i - \mu)(X_i - \mu)^T$$


ITS STILL JUST COUNTING!

22 Sep 2011

11755/18797

20

Laplacian



$$P(x) = L(x; \mu, b) = \frac{1}{2b} \exp\left(-\frac{|x - \mu|}{b}\right)$$

- Parameters: Mean μ , scale b ($b > 0$)

22 Sep 2011

11755/18797

21

Maximum Likelihood: Laplacian

- Given a collection of observations (x_1, x_2, \dots) , estimate mean μ and scale b

$$\log(P(x_1, x_2, \dots)) = C - N \log(b) - \sum_i \frac{|x_i - \mu|}{b}$$

- Maximizing w.r.t μ and b gives us

$$\mu = \frac{1}{N} \sum_i x_i \quad b = \frac{1}{N} \sum_i |x_i - \mu|$$

22 Sep 2011

11755/18797

22

Dirichlet

(from wikipedia)





Fig 3. Clockwise from top left: $\alpha=(6, 2, 2)$, $(3, 7, 9)$, $(6, 2, 6)$, $(2, 3, 4)$



log of the density as we change α from $\alpha=(0.3, 0.3, 0.3)$ to $(2.0, 2.0, 2.0)$, keeping all the individual α 's equal to each other.

$$P(X) = D(X; \alpha) = \frac{\prod_i \Gamma(\alpha_i)}{\Gamma(\sum_i \alpha_i)} \prod_i x_i^{\alpha_i - 1}$$

- Parameters are α s
 - Determine mode and curvature
- Defined only of probability vectors
 - $X = [x_1 \ x_2 \ \dots \ x_K]$, $\sum_i x_i = 1$, $x_i \geq 0$ for all i

22 Sep 2011

11755/18797

23

Maximum Likelihood: Dirichlet

- Given a collection of observations (X_1, X_2, \dots) , estimate α

$$\log(P(X_1, X_2, \dots)) = \sum_i \sum_j (\alpha_j - 1) \log(X_{j,i}) + N \sum_j \log(\Gamma(\alpha_j)) - N \log\left(\Gamma\left(\sum_i \alpha_i\right)\right)$$


- No closed form solution for α s.
 - Needs gradient ascent
- Several distributions have this property: the ML estimate of their parameters have no closed form solution

22 Sep 2011

11755/18797

24

Continuing the Thought Experiment



6 3 1 5 4 1 2 4 ... 4 4 1 6 3 2 1 2 ...


- Two persons shoot loaded dice repeatedly
 - The dice are differently loaded for the two of them
- We observe the series of outcomes for both persons

How to determine the probability distributions of the two dice?

22 Sep 2011 11755/18797 25

Estimating Probabilities


- Observation: The sequence of numbers from the two dice
 - As indicated by the colors, we know who rolled what number



22 Sep 2011 11755/18797 26

Estimating Probabilities

- Observation: The sequence of numbers from the two dice
 - As indicated by the colors, we know who rolled what number
- Segregation: Separate the blue observations from the red




22 Sep 2011 11755/18797 27

Estimating Probabilities


- Observation: The sequence of numbers from the two dice
 - As indicated by the colors, we know who rolled what number
- Segregation: Separate the blue observations from the red
- From each set compute probabilities for each of the 6 possible outcomes

$P(\text{number}) = \frac{\text{no. of times number was rolled}}{\text{total number of observed rolls}}$



22 Sep 2011 11755/18797 28

A Thought Experiment




6 3 1 5 4 1 2 4 ... 4 4 1 6 3 2 1 2 ...

6 4 1 5 3 2 2 2 ...

- Now imagine that you cannot observe the dice yourself
- Instead there is a "caller" who randomly calls out the outcomes
 - 40% of the time he calls out the number from the left shooter, and 60% of the time, the one from the right (and you know this)
- At any time, you do not know which of the two he is calling out
- How do you determine the probability distributions for the two dice?

22 Sep 2011 11755/18797 29

A Thought Experiment



6 3 1 5 4 1 2 4 ... 4 4 1 6 3 2 1 2 ...

6 4 1 5 3 2 2 2 ...

- How do you now determine the probability distributions for the two sets of dice ...
- .. If you do not even know what fraction of time the blue numbers are called, and what fraction are red?

22 Sep 2011 11755/18797 30

A Mixture Multinomial

- The caller will call out a number X in any given callout IF
 - He selects "RED", and the Red die rolls the number X
 - OR
 - He selects "BLUE" and the Blue die rolls the number X
- $P(X) = P(\text{Red})P(X|\text{Red}) + P(\text{Blue})P(X|\text{Blue})$
 - E.g. $P(6) = P(\text{Red})P(6|\text{Red}) + P(\text{Blue})P(6|\text{Blue})$
- A distribution that *combines* (or *mixes*) multiple multinomials is a *mixture* multinomial

$$P(X) = \sum_Z P(Z)P(X|Z)$$

Mixture weights Component multinomials

22 Sep 2011

11755/18797

31

Mixture Distributions

$$P(X) = \sum_Z P(Z)P(X|Z) \quad \text{Mixture Gaussian} \quad P(X) = \sum_Z P(Z)N(X; \mu_z, \Theta_z)$$

Mixture weights Component distributions

Mixture of Gaussians and Laplacians

$$P(X) = \sum_Z P(Z)N(X; \mu_z, \Theta_z) + \sum_Z P(Z) \prod_i L(X_i; \mu_z, b_{z,i})$$

- Mixture distributions mix several component distributions
 - Component distributions may be of varied type
- Mixing weights must sum to 1.0
- Component distributions integrate to 1.0
- Mixture distribution integrates to 1.0

22 Sep 2011

11755/18797

32

Maximum Likelihood Estimation

- For our problem: $P(X) = \sum_Z P(Z)P(X|Z)$
 - Z = color of dice
- $P(n_1, n_2, n_3, n_4, n_5, n_6) = \text{Const} \prod_X P(X)^{n_X} = \text{Const} \prod_X \left(\sum_Z P(Z)P(X|Z) \right)^{n_X}$
- Maximum likelihood solution: Maximize
 - $\log(P(n_1, n_2, n_3, n_4, n_5, n_6)) = \log(\text{Const}) + \sum_X n_X \log \left(\sum_Z P(Z)P(X|Z) \right)$
- No closed form solution (summation inside log)!
 - In general ML estimates for mixtures do not have a closed form
 - USE EM!

22 Sep 2011

11755/18797

33

Expectation Maximization

- It is possible to estimate all parameters in this setup using the Expectation Maximization (or EM) algorithm
- First described in a landmark paper by Dempster, Laird and Rubin
 - Maximum Likelihood Estimation from incomplete data, via the EM Algorithm, Journal of the Royal Statistical Society, Series B, 1977
- Much work on the algorithm since then
- The principles behind the algorithm existed for several years prior to the landmark paper, however.

22 Sep 2011

11755/18797

34

Expectation Maximization

- Iterative solution
- Get some initial estimates for all parameters
 - Dice shooter example: This includes probability distributions for dice AND the probability with which the caller selects the dice
- Two steps that are iterated:
 - **Expectation Step:** Estimate statistically, the values of *unseen* variables
 - **Maximization Step:** Using the estimated values of the unseen variables as truth, estimates of the model parameters

22 Sep 2011

11755/18797

35

EM: The auxiliary function

- EM iteratively optimizes the following auxiliary function
- $Q(\theta, \theta') = \sum_Z P(Z|X, \theta') \log(P(Z, X | \theta))$
 - Z are the unseen variables
 - Assuming Z is discrete (may not be)
- θ' are the parameter estimates from the previous iteration
- θ are the estimates to be obtained in the current iteration

22 Sep 2011

11755/18797

36

Expectation Maximization as counting

- Hidden variable: Z
 - Dice: The identity of the dice whose number has been called out
- If we knew Z for every observation, we could estimate all terms
 - By adding the observation to the right bin
- Unfortunately, we do not know Z – it is hidden from us!
- Solution: FRAGMENT THE OBSERVATION

22 Sep 2011 11755/18797 37

Fragmenting the Observation

- EM is an iterative algorithm
 - At each time there is a *current* estimate of parameters
- The “size” of the fragments is proportional to the *a posteriori probability* of the component distributions
 - The *a posteriori* probabilities of the various values of Z are computed using Bayes' rule:

$$P(Z | X) = \frac{P(X | Z)P(Z)}{P(X)} = CP(X | Z)P(Z)$$

- Every dice gets a fragment of size $P(\text{dice} | \text{number})$

22 Sep 2011 11755/18797 38

Expectation Maximization

- Hypothetical Dice Shooter Example:
- We obtain an initial estimate for the probability distribution of the two sets of dice (somehow):

- We obtain an initial estimate for the probability with which the caller calls out the two shooters (somehow)

22 Sep 2011 11755/18797 39

Expectation Maximization

- Hypothetical Dice Shooter Example:
- Initial estimate:
 - $P(\text{blue}) = P(\text{red}) = 0.5$
 - $P(4 | \text{blue}) = 0.1$, for $P(4 | \text{red}) = 0.05$
- Caller has just called out 4
- Posterior probability of colors:
 - $P(\text{red} | X = 4) = CP(X = 4 | Z = \text{red})P(Z = \text{red}) = C \times 0.05 \times 0.5 = C \times 0.025$
 - $P(\text{blue} | X = 4) = CP(X = 4 | Z = \text{blue})P(Z = \text{blue}) = C \times 0.1 \times 0.5 = C \times 0.05$

Normalizing : $P(\text{red} | X = 4) = 0.33$; $P(\text{blue} | X = 4) = 0.67$

22 Sep 2011 11755/18797 40

Expectation Maximization

22 Sep 2011 11755/18797 41

Expectation Maximization

- Every observed roll of the dice contributes to both “Red” and “Blue”

22 Sep 2011 11755/18797 42

Expectation Maximization

- Every observed roll of the dice contributes to both "Red" and "Blue"

22 Sep 2011 11755/18797 43

Expectation Maximization

- Every observed roll of the dice contributes to both "Red" and "Blue"

22 Sep 2011 11755/18797 44

Expectation Maximization

- Every observed roll of the dice contributes to both "Red" and "Blue"

22 Sep 2011 11755/18797 45

Expectation Maximization

- Every observed roll of the dice contributes to both "Red" and "Blue"

22 Sep 2011 11755/18797 46

Expectation Maximization

- Every observed roll of the dice contributes to both "Red" and "Blue"
- Total count for "Red" is the sum of all the posterior probabilities in the red column
 - 7.31
- Total count for "Blue" is the sum of all the posterior probabilities in the blue column
 - 10.69
- Note: $10.69 + 7.31 = 18$ = the total number of instances

Called	P(red X)	P(blue X)
6	.8	.2
4	.33	.67
5	.33	.67
1	.57	.43
2	.14	.86
3	.33	.67
4	.33	.67
5	.33	.67
2	.14	.86
2	.14	.86
1	.57	.43
4	.33	.67
3	.33	.67
4	.33	.67
6	.8	.2
2	.14	.86
1	.57	.43
6	.8	.2

7.31 10.69

22 Sep 2011 11755/18797 47

Expectation Maximization

- Total count for "Red" : 7.31
- Red:
 - Total count for 1: 1.71

Called	P(red X)	P(blue X)
6	.8	.2
4	.33	.67
5	.33	.67
1	.57	.43
2	.14	.86
3	.33	.67
4	.33	.67
5	.33	.67
2	.14	.86
2	.14	.86
1	.57	.43
4	.33	.67
3	.33	.67
4	.33	.67
6	.8	.2
2	.14	.86
1	.57	.43
6	.8	.2

7.31 10.69

22 Sep 2011 11755/18797 48

Expectation Maximization

- Total count for "Red" : 7.31
- Red:
 - Total count for 1: 1.71
 - Total count for 2: 0.56

Called	P(red X)	P(blue X)
6	.8	.2
4	.33	.67
5	.33	.67
1	.57	.43
2	.14	.86
3	.33	.67
4	.33	.67
5	.33	.67
2	.14	.86
2	.14	.86
1	.57	.43
4	.33	.67
3	.33	.67
4	.33	.67
6	.8	.2
2	.14	.86
1	.57	.43
6	.8	.2
7.31		10.69

22 Sep 2011

11755/18797

49

Expectation Maximization

- Total count for "Red" : 7.31
- Red:
 - Total count for 1: 1.71
 - Total count for 2: 0.56
 - Total count for 3: 0.66

Called	P(red X)	P(blue X)
6	.8	.2
4	.33	.67
5	.33	.67
1	.57	.43
2	.14	.86
3	.33	.67
4	.33	.67
5	.33	.67
2	.14	.86
2	.14	.86
1	.57	.43
4	.33	.67
3	.33	.67
4	.33	.67
6	.8	.2
2	.14	.86
1	.57	.43
6	.8	.2
7.31		10.69

22 Sep 2011

11755/18797

50

Expectation Maximization

- Total count for "Red" : 7.31
- Red:
 - Total count for 1: 1.71
 - Total count for 2: 0.56
 - Total count for 3: 0.66
 - Total count for 4: 1.32

Called	P(red X)	P(blue X)
6	.8	.2
4	.33	.67
5	.33	.67
1	.57	.43
2	.14	.86
3	.33	.67
4	.33	.67
5	.33	.67
2	.14	.86
2	.14	.86
1	.57	.43
4	.33	.67
3	.33	.67
4	.33	.67
6	.8	.2
2	.14	.86
1	.57	.43
6	.8	.2
7.31		10.69

22 Sep 2011

11755/18797

51

Expectation Maximization

- Total count for "Red" : 7.31
- Red:
 - Total count for 1: 1.71
 - Total count for 2: 0.56
 - Total count for 3: 0.66
 - Total count for 4: 1.32
 - Total count for 5: 0.66

Called	P(red X)	P(blue X)
6	.8	.2
4	.33	.67
5	.33	.67
1	.57	.43
2	.14	.86
3	.33	.67
4	.33	.67
5	.33	.67
2	.14	.86
2	.14	.86
1	.57	.43
4	.33	.67
3	.33	.67
4	.33	.67
6	.8	.2
2	.14	.86
1	.57	.43
6	.8	.2
7.31		10.69

22 Sep 2011

11755/18797

52

Expectation Maximization

- Total count for "Red" : 7.31
- Red:
 - Total count for 1: 1.71
 - Total count for 2: 0.56
 - Total count for 3: 0.66
 - Total count for 4: 1.32
 - Total count for 5: 0.66
 - Total count for 6: 2.4

Called	P(red X)	P(blue X)
6	.8	.2
4	.33	.67
5	.33	.67
1	.57	.43
2	.14	.86
3	.33	.67
4	.33	.67
5	.33	.67
2	.14	.86
2	.14	.86
1	.57	.43
4	.33	.67
3	.33	.67
4	.33	.67
6	.8	.2
2	.14	.86
1	.57	.43
6	.8	.2
7.31		10.69

22 Sep 2011

11755/18797

53

Expectation Maximization

- Total count for "Red" : 7.31
- Red:
 - Total count for 1: 1.71
 - Total count for 2: 0.56
 - Total count for 3: 0.66
 - Total count for 4: 1.32
 - Total count for 5: 0.66
 - Total count for 6: 2.4
- Updated probability of Red dice:
 - $P(1 | \text{Red}) = 1.71/7.31 = 0.234$
 - $P(2 | \text{Red}) = 0.56/7.31 = 0.077$
 - $P(3 | \text{Red}) = 0.66/7.31 = 0.090$
 - $P(4 | \text{Red}) = 1.32/7.31 = 0.181$
 - $P(5 | \text{Red}) = 0.66/7.31 = 0.090$
 - $P(6 | \text{Red}) = 2.40/7.31 = 0.328$

Called	P(red X)	P(blue X)
6	.8	.2
4	.33	.67
5	.33	.67
1	.57	.43
2	.14	.86
3	.33	.67
4	.33	.67
5	.33	.67
2	.14	.86
2	.14	.86
1	.57	.43
4	.33	.67
3	.33	.67
4	.33	.67
6	.8	.2
2	.14	.86
1	.57	.43
6	.8	.2
7.31		10.69

22 Sep 2011

11755/18797

54

Expectation Maximization

- Total count for "Blue" : 10.69
- Blue:
 - Total count for 1: 1.29

Called	P(red X)	P(blue X)
6	.8	.2
4	.33	.67
5	.33	.67
1	.57	.43
2	.14	.86
3	.33	.67
4	.33	.67
5	.33	.67
2	.14	.86
2	.14	.86
1	.57	.43
4	.33	.67
3	.33	.67
4	.33	.67
6	.8	.2
2	.14	.86
1	.57	.43
6	.8	.2

7.31 10.69

22 Sep 2011

11755/18797

55

Expectation Maximization

- Total count for "Blue" : 10.69
- Blue:
 - Total count for 1: 1.29
 - Total count for 2: 3.44

Called	P(red X)	P(blue X)
6	.8	.2
4	.33	.67
5	.33	.67
1	.57	.43
2	.14	.86
3	.33	.67
4	.33	.67
5	.33	.67
2	.14	.86
2	.14	.86
1	.57	.43
4	.33	.67
3	.33	.67
4	.33	.67
6	.8	.2
2	.14	.86
1	.57	.43
6	.8	.2

7.31 10.69

22 Sep 2011

11755/18797

56

Expectation Maximization

- Total count for "Blue" : 10.69
- Blue:
 - Total count for 1: 1.29
 - Total count for 2: 3.44
 - Total count for 3: 1.34

Called	P(red X)	P(blue X)
6	.8	.2
4	.33	.67
5	.33	.67
1	.57	.43
2	.14	.86
3	.33	.67
4	.33	.67
5	.33	.67
2	.14	.86
2	.14	.86
1	.57	.43
4	.33	.67
3	.33	.67
4	.33	.67
6	.8	.2
2	.14	.86
1	.57	.43
6	.8	.2

7.31 10.69

22 Sep 2011

11755/18797

57

Expectation Maximization

- Total count for "Blue" : 10.69
- Blue:
 - Total count for 1: 1.29
 - Total count for 2: 3.44
 - Total count for 3: 1.34
 - Total count for 4: 2.68

Called	P(red X)	P(blue X)
6	.8	.2
4	.33	.67
5	.33	.67
1	.57	.43
2	.14	.86
3	.33	.67
4	.33	.67
5	.33	.67
2	.14	.86
2	.14	.86
1	.57	.43
4	.33	.67
3	.33	.67
4	.33	.67
6	.8	.2
2	.14	.86
1	.57	.43
6	.8	.2

7.31 10.69

22 Sep 2011

11755/18797

58

Expectation Maximization

- Total count for "Blue" : 10.69
- Blue:
 - Total count for 1: 1.29
 - Total count for 2: 3.44
 - Total count for 3: 1.34
 - Total count for 4: 2.68
 - Total count for 5: 1.34

Called	P(red X)	P(blue X)
6	.8	.2
4	.33	.67
5	.33	.67
1	.57	.43
2	.14	.86
3	.33	.67
4	.33	.67
5	.33	.67
2	.14	.86
2	.14	.86
1	.57	.43
4	.33	.67
3	.33	.67
4	.33	.67
6	.8	.2
2	.14	.86
1	.57	.43
6	.8	.2

7.31 10.69

22 Sep 2011

11755/18797

59

Expectation Maximization

- Total count for "Blue" : 10.69
- Blue:
 - Total count for 1: 1.29
 - Total count for 2: 3.44
 - Total count for 3: 1.34
 - Total count for 4: 2.68
 - Total count for 5: 1.34
 - Total count for 6: 0.6

Called	P(red X)	P(blue X)
6	.8	.2
4	.33	.67
5	.33	.67
1	.57	.43
2	.14	.86
3	.33	.67
4	.33	.67
5	.33	.67
2	.14	.86
2	.14	.86
1	.57	.43
4	.33	.67
3	.33	.67
4	.33	.67
6	.8	.2
2	.14	.86
1	.57	.43
6	.8	.2

7.31 10.69

22 Sep 2011

11755/18797

60

Expectation Maximization

- Total count for "Blue" : 10.69
- Blue:
 - Total count for 1: 1.29
 - Total count for 2: 3.44
 - Total count for 3: 1.34
 - Total count for 4: 2.68
 - Total count for 5: 1.34
 - Total count for 6: 0.6

Updated probability of Blue dice:

- P(1 | Blue) = 1.29/11.69 = 0.122
- P(2 | Blue) = 0.56/11.69 = 0.322
- P(3 | Blue) = 0.66/11.69 = 0.125
- P(4 | Blue) = 1.32/11.69 = 0.250
- P(5 | Blue) = 0.66/11.69 = 0.125
- P(6 | Blue) = 2.40/11.69 = 0.056

Called	P(red X)	P(blue X)
6	.8	.2
4	.33	.67
5	.33	.67
1	.57	.43
2	.14	.86
3	.33	.67
4	.33	.67
5	.33	.67
2	.14	.86
2	.14	.86
1	.57	.43
4	.33	.67
3	.33	.67
4	.33	.67
6	.8	.2
2	.14	.86
1	.57	.43
6	.8	.2

7.31 10.69

22 Sep 2011

11755/18797

61

Expectation Maximization

- Total count for "Red" : 7.31
- Total count for "Blue" : 10.69
- Total instances = 18
 - Note 7.31+10.69 = 18
- We also revise our estimate for the probability that the caller calls out Red or Blue
 - i.e the fraction of times that he calls Red and the fraction of times he calls Blue

- P(Z=Red) = 7.31/18 = 0.41
- P(Z=Blue) = 10.69/18 = 0.59

Called	P(red X)	P(blue X)
6	.8	.2
4	.33	.67
5	.33	.67
1	.57	.43
2	.14	.86
3	.33	.67
4	.33	.67
5	.33	.67
2	.14	.86
2	.14	.86
1	.57	.43
4	.33	.67
3	.33	.67
4	.33	.67
6	.8	.2
2	.14	.86
1	.57	.43
6	.8	.2

7.31 10.69

22 Sep 2011

11755/18797

62

The updated values

Probability of Red dice:

- P(1 | Red) = 1.71/7.31 = 0.234
- P(2 | Red) = 0.56/7.31 = 0.077
- P(3 | Red) = 0.66/7.31 = 0.090
- P(4 | Red) = 1.32/7.31 = 0.181
- P(5 | Red) = 0.66/7.31 = 0.090
- P(6 | Red) = 2.40/7.31 = 0.328

Probability of Blue dice:

- P(1 | Blue) = 1.29/11.69 = 0.122
- P(2 | Blue) = 0.56/11.69 = 0.322
- P(3 | Blue) = 0.66/11.69 = 0.125
- P(4 | Blue) = 1.32/11.69 = 0.250
- P(5 | Blue) = 0.66/11.69 = 0.125
- P(6 | Blue) = 2.40/11.69 = 0.056

- P(Z=Red) = 7.31/18 = 0.41
- P(Z=Blue) = 10.69/18 = 0.59

THE UPDATED VALUES CAN BE USED TO REPEAT THE PROCESS. ESTIMATION IS AN ITERATIVE PROCESS

Called	P(red X)	P(blue X)
6	.8	.2
4	.33	.67
5	.33	.67
1	.57	.43
2	.14	.86
3	.33	.67
4	.33	.67
5	.33	.67
2	.14	.86
2	.14	.86
1	.57	.43
4	.33	.67
3	.33	.67
4	.33	.67
6	.8	.2
2	.14	.86
1	.57	.43
6	.8	.2

22 Sep 2011

11755/18797

63

The Dice Shooter Example



- Initialize $P(Z)$, $P(X|Z)$
- Estimate $P(Z|X)$ for each Z , for each called out number
 - Associate X with each value of Z , with weight $P(Z|X)$
- Re-estimate $P(X|Z)$ for every value of X and Z
- Re-estimate $P(Z)$
- If not converged, return to 2

22 Sep 2011

11755/18797

64

In Squiggles

- Given a sequence of observations O_1, O_2, \dots
 - N_x is the number of observations of number X
- Initialize $P(Z)$, $P(X|Z)$ for dice Z and numbers X

Iterate:

- For each number X :
$$P(Z|X) = \frac{P(X|Z)P(Z)}{\sum_{Z'} P(Z')P(X|Z')}$$

Update:

$$P(X|Z) = \frac{\sum_{O \text{ such that } O=X} P(Z|O)}{\sum_O P(Z|O)} = \frac{N_X P(Z|X)}{\sum_X N_X P(Z|X)}$$

$$P(Z) = \frac{\sum_X N_X P(Z|X)}{\sum_{Z'} \sum_X N_X P(Z'|X)}$$

22 Sep 2011

11755/18797

65

Solutions may not be unique

- The EM algorithm will give us one of many solutions, all equally valid!

- The probability of 6 being called out:

$$P(6) = \alpha P(6|red) + \beta P(6|blue) = \alpha P_r + \beta P_b$$

- Assigns P_r as the probability of 6 for the red die
- Assigns P_b as the probability of 6 for the blue die

- The following too is a valid solution [FIX]

$$P(6) = 1.0(\alpha P_r + \beta P_b) + 0.0 \text{ anything}$$

- Assigns 1.0 as the a priori probability of the red die
- Assigns 0.0 as the probability of the blue die


- The solution is NOT unique

22 Sep 2011

11755/18797

66

A More Complex Model




$$P(X) = \sum_k P(k)N(X; \mu_k, \Theta_k) = \sum_k \frac{P(k)}{\sqrt{(2\pi)^d |\Theta_k|}} \exp(-0.5(X - \mu_k)^T \Theta_k^{-1}(X - \mu_k))$$

- Gaussian mixtures are often good models for the distribution of multivariate data
- Problem: Estimating the parameters, given a collection of data

22 Sep 2011 11755/18797 67

Gaussian Mixtures: Generating model

6.1 1.4 5.3 1.9 4.2 2.2 4.9 0.5


$$P(X) = \sum_k P(k)N(X; \mu_k, \Theta_k)$$


- The caller now has two Gaussians
 - At each draw he randomly selects a Gaussian, by the mixture weight distribution
 - He then draws an observation from that Gaussian
 - Much like the dice problem (only the outcomes are now real numbers and can be anything)

22 Sep 2011 11755/18797 68

Estimating GMM with complete information

- Observation: A collection of numbers drawn from a mixture of 2 Gaussians
 - As indicated by the colors, we know which Gaussian generated what number
- Segregation: Separate the blue observations from the red
- From each set compute parameters for that Gaussian




$$\mu_{red} = \frac{1}{N_{red}} \sum_{i \in red} X_i \quad \Theta_{red} = \frac{1}{N_{red}} \sum_{i \in red} (X_i - \mu_{red})(X_i - \mu_{red})^T$$

$$P(red) = \frac{N_{red}}{N}$$

22 Sep 2011 11755/18797 69

Fragmenting the observation



- The identity of the Gaussian is not known!
- Solution: **Fragment the observation**
- Fragment size proportional to a *posteriori* probability

$$P(k | X) = \frac{P(X | k)P(k)}{\sum_{k'} P(k')P(X | k')} = \frac{P(k)N(X; \mu_k, \Theta_k)}{\sum_{k'} P(k')N(X; \mu_{k'}, \Theta_{k'})}$$

22 Sep 2011 11755/18797 70

Expectation Maximization

- Initialize $P(k)$, μ_k and Θ_k for both Gaussians
 - Important how we do this
 - Typical solution: Initialize means randomly, Θ_k as the global covariance of the data and $P(k)$ uniformly
- Compute fragment sizes for each Gaussian, for each observation

Number	P(red X)	P(blue X)
6.1	.81	.19
1.4	.33	.67
5.3	.75	.25
1.9	.41	.59
4.2	.64	.36
2.2	.43	.57
4.9	.66	.34
0.5	.05	.95

$$P(k | X) = \frac{P(k)N(X; \mu_k, \Theta_k)}{\sum_{k'} P(k')N(X; \mu_{k'}, \Theta_{k'})}$$

22 Sep 2011 11755/18797 71

Expectation Maximization

- Each observation contributes only as much as its fragment size to each statistic
- Mean(red) = $(6.1 \cdot 0.81 + 1.4 \cdot 0.33 + 5.3 \cdot 0.75 + 1.9 \cdot 0.41 + 4.2 \cdot 0.64 + 2.2 \cdot 0.43 + 4.9 \cdot 0.66 + 0.5 \cdot 0.05) / (0.81 + 0.33 + 0.75 + 0.41 + 0.64 + 0.43 + 0.66 + 0.05) = 17.05 / 4.08 = 4.18$
- Var(red) = $((6.1-4.18)^2 \cdot 0.81 + (1.4-4.18)^2 \cdot 0.33 + (5.3-4.18)^2 \cdot 0.75 + (1.9-4.18)^2 \cdot 0.41 + (4.2-4.18)^2 \cdot 0.64 + (2.2-4.18)^2 \cdot 0.43 + (4.9-4.18)^2 \cdot 0.66 + (0.5-4.18)^2 \cdot 0.05) / (0.81 + 0.33 + 0.75 + 0.41 + 0.64 + 0.43 + 0.66 + 0.05)$

Number	P(red X)	P(blue X)
6.1	.81	.19
1.4	.33	.67
5.3	.75	.25
1.9	.41	.59
4.2	.64	.36
2.2	.43	.57
4.9	.66	.34
0.5	.05	.95

$$P(red) = \frac{4.08}{8}$$

22 Sep 2011 11755/18797 72

EM for Gaussian Mixtures

1. Initialize $P(k)$, μ_k and Θ_k for all Gaussians
2. For each observation X compute a *posteriori* probabilities for all Gaussian

$$P(k | X) = \frac{P(k)N(X; \mu_k, \Theta_k)}{\sum_{k'} P(k')N(X; \mu_{k'}, \Theta_{k'})}$$

3. Update mixture weights, means and variances for all Gaussians

$$P(k) = \frac{\sum_x P(k|x)}{N} \quad \mu_k = \frac{\sum_x P(k|x) X}{\sum_x P(k|x)} \quad \Theta_k = \frac{\sum_x P(k|x) (X - \mu_k)^2}{\sum_x P(k|x)}$$

4. If not converged, return to 2


22 Sep 2011

11755/18797

73

EM estimation of Gaussian Mixtures

- An Example



Histogram of 4000 instances of a randomly generated data

Individual parameters of a two-Gaussian mixture estimated by EM

Two-Gaussian mixture estimated by EM

22 Sep 2011

11755/18797

74

Expectation Maximization

- The same principle can be extended to mixtures of other distributions.
- E.g. Mixture of Laplacians: Laplacian parameters become

$$\mu_k = \frac{1}{\sum_x P(k|x)} \sum_x P(k|x)x \quad b_k = \frac{1}{\sum_x P(k|x)} \sum_x P(k|x) |x - \mu_k|$$

- In a mixture of Gaussians and Laplacians, Gaussians use the Gaussian update rules, Laplacians use the Laplacian rule

22 Sep 2011

11755/18797

75

Expectation Maximization

- The EM algorithm is used whenever proper statistical analysis of a phenomenon requires the knowledge of a hidden or missing variable (or a set of hidden/missing variables)
 - The hidden variable is often called a "latent" variable
- Some examples:
 - Estimating mixtures of distributions
 - Only data are observed. The individual distributions and mixing proportions must both be learnt.
 - Estimating the distribution of data, when some attributes are missing
 - Estimating the dynamics of a system, based only on observations that may be a complex function of system state

22 Sep 2011

11755/18797

76

Solve this problem:


- Caller rolls a dice and flips a coin
- He calls out the number rolled if the coin shows head
- Otherwise he calls the number+1
- Determine $p(\text{heads})$ and $p(\text{number})$ for the dice from a collection of outputs
- Caller rolls two dice
- He calls out the sum
- Determine $P(\text{dice})$ from a collection of outputs

22 Sep 2011

11755/18797

77

The dice and the coin




- Unknown: Whether it was head or tails

22 Sep 2011

11755/18797

78

The two dice



- Unknown: How to partition the number
- $\text{Count}_{\text{blue}}(3) \propto P(3,1 | 4)$
- $\text{Count}_{\text{blue}}(2) \propto P(2,2 | 4)$
- $\text{Count}_{\text{blue}}(1) \propto P(1,3 | 4)$


22 Sep 2011

11755/18797

79

Fragmentation can be hierarchical

$$P(X) = \sum_k P(k) \sum_Z P(Z | k) P(X | Z, k)$$



- E.g. mixture of mixtures
- Fragments are further fragmented..
 - Work this out

22 Sep 2011

11755/18797

80

More later

- Will see a couple of other instances of the use of EM
- Work out HMM training
 - Assume state output distributions are multinomials
 - Assume they are Gaussian
 - Assume Gaussian mixtures

22 Sep 2011

11755/18797

81